



Seen by me, Suzanne Beck, candidate-civil law notary in Rotterdam, the Netherlands, as a substitute of Victor Joseph Antonius Johannes Clemens van Heeswijk, civil law notary in Rotterdam, The Netherlands, for legalisation of the signatures placed on the attached document of:

- mr Christiaan Beindorff, c/o Unilever Research Vlaardingen, Olivier van Noortlaan 120, 3133 AT Vlaardingen, the Netherlands;
- mr Frederick William Cain, residing at dr. Blookerstraat 12, 2271 VL Voorburg, the Netherlands;
- mr John Hugh Pierce, residing at 24, Purbeck Road, Rushden, Northants, NN10 9UH, Great Britain;
- mrs Ulrike Schmid, c/o Loders Croklaan B.V., Hogeweg 1, 1521 AZ Wormerveer, the Netherlands;
- mr Erik Schweitzer, c/o Loders Croklaan B.V., Hogeweg 1, 1521 AZ Wormerveer, the Netherlands;
- mr Jeroen Nicolaas M. van Straalen, residing at Kerkstraat 94, 1687 AT Wognum, the Netherlands.

Rotterdam, 12 July 2001



A large, stylized handwritten signature, "Beck", is written in black ink. The signature is enclosed within a large, loopy oval shape. A long horizontal line extends from the bottom of the oval.

APOSTILLE

(Convention de La Haye du 5 Octobre 1961)

1. Country: The NETHERLANDS

This public document

2. has been signed by mr S.

Beck

3. acting in the capacity of

candidate notary

4. bears the stamp of

mr V.J.A.J.C. van Heeswijk

Certified

5. at ROTTERDAM 6. on 12/07/2001



7. by the clerk of the District Court

(Griffier van de Arrondissementsrechtbank)

8. N° 109900

9. Stamp:

10. Signature:





Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

**Blatt 2 der Bescheinigung
Sheet 2 of the certificate
Page 2 de l'attestation**

Anmeldung Nr.:
Application no.:
Demande n°: 00304753.7

Anmeldetag:
Date of filing: 05/06/00
Date de dépôt:

Anmelder:
Applicant(s):
Demandeur(s):
UNILEVER PLC
London EC4P 4BQ
UNITED KINGDOM

Bezeichnung der Erfindung:
Title of the invention:
Titre de l'invention:
Blends of ursolic acid/oleanolic acid

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:
State:
Pays:

Tag:
Date:
Date:

Aktenzeichen:
File no.
Numéro de dépôt:

Internationale Patentklassifikation:
International Patent classification:
Classification internationale des brevets:

C07J63/00, A61K35/78, A23D9/007, A23D7/00

Am Anmeldetag benannte Vertragsstaaten:
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE/UK
Etats contractants désignés lors du dépôt:

Bemerkungen:
Remarks:
Remarques:

THIS PAGE BLANK (USPTO)

BLENDS OF URSOLIC ACID/OLEANOLIC ACID

Ursolic acid and oleanolic acid are known compounds that are also known for their health effects; references

5 wherein these effects are disclosed are eg JP 09/040 689; JP 09/ 067 249; CN 1 085 748; JP 1 039 973; JP 03/287 531; JP 03/287 430; EP 774 255; JP 07/258 098; JP 07/048 260; JP 01/132 531; FR 2 535 203; and JP 1-207 262.

Compositions comprising ursolic acid and oleanolic acid in
10 weight ratios of 1:99 to 99:1 can be obtained by extracting natural sources like fruit skins or herbs, in particular by extracting skins of apples, pears, cranberries, cherries and prunes. The extracts so obtained contain ursolic acid and oleanolic acid in amounts of about 5-60 % and in weight
15 ratios mentioned above. However these extracts have a major drawback ie the extracts display a severe negative off flavour and as the components are intended to be used as health components in food products and the consumer does not appreciate a negative off flavour while consuming his
20 food it would be a main advantage when mixtures of these components could be obtained containing sufficient amounts of ursolic acid and oleanolic acid to make them useful for application in foods as health component and that do not display the negative off flavour of the products available
25 so far.

We studied whether we could obtain such products. In this study we found that the negative off flavour of the ursolic acid/oleanolic acid mixture is due to the presence of
30 natural apolar and/or low molecular weight components in the natural extracts known so far. The apolar components being defined as that fraction of an extract from fruit

skins that is eluted from a polar silica gel column, preferably with a Alltech Econoshpere Silica HPLC column (150*4.6 mm, 5 μ m) with an eluent, preferably being either a mixture of hexane/toluene (50:50) or a mixture of 5 toluene/ethylacetate/formic acid (500/200/16), in 0-7 min. Low molecular weight components are detected by GC with a apolar column, preferably a Chrompack SIMDIST (10m*0.53mm, 0.1 μ m filmthickness) or a Quadrex DB-5 (10m*0.53mm, 0.1 μ m filmthickness) and a temperature program, preferably 10 starting with 120 oC (1 min); then 20oC/min until 325 oC, then 5 oC/min until 350 oC (5 min) with retention times from 0.5-7 min. The preferably used carrier gas is hydrogen with a pressure of 15 psi.

15 We found a method wherewith the amount of these apolar and/or low molecular weight components could be reduced considerably i.e. to below a level that would result in a negative off flavour impression. The result of this method is the first part of our invention i.e. a mixture 20 comprising ursolic acid and oleanolic acid in a weight ratio of 1:99 to 99:1, preferably 5:95 to 95:5, most preferably 15:85 to 15:85 wherein the mixture contains less than 20 wt %, preferably less than 10 wt %, most preferably 1 to 6 wt % of the natural apolar and/or low molecular 25 weight components as present in natural extracts for ursolic acid and oleanolic acid. The apolar and/or low molecular weight components were found to be components belonging to the class of hydrocarbons, alcohols, fatty acids, triglycerides, ketones and carbohydrates

30

Although the above mix could be used as such in food products it is often preferred to use it as a blend with

other components in particular as a blend with glycerides, preferably triglycerides. Therefore according to another embodiment of our invention we also found blends of a health component and a glyceride wherein the health
5 component is the mixture according to the invention disclosed above and which blend contains 1 to 99 wt % of one or more components selected from mono-, di-, and triglycerides as the glyceride.

The glyceride part of this blend preferably displays a
10 solid fat content measured by NMR-pulse on a non-stabilised fat at the temperature indicated of :

5 to 90 at 5 °C

2 to 80 at 20 °C and

less than 15, preferably less than 10 at 35 °C.

15 The solid fat content is measured by the well known NMR - pulse technique on a fat that is not stabilised, this means that the measurement was performed on a fat that was subjected to the following treatment:

Melt at 80 °C, keep it at 80 °C for 15 min, cool it to 0 °C
20 and keep it at 0 °C for 30 min, heat it to measurement temperature and keep it thereon for 30 min and measure the N-value at this temperature.

Preferred blends that are good applicable in food products
25 are blends comprising components A, B and C, wherein:

A = the health composition according to the invention and as formulated in claims 1 or 2

B = a solid fat with an N20 of more than 20 , preferably more than 45 , most preferably more than 60 and

30 C - a fat having at least 40 wt % of fatty acids with 18 C-atoms and having one to three double bonds

A being present in amounts of more than 0.1 wt %, preferably 0.1 to 20 wt %, most preferably 0.2 to 10 wt %
B being present in amounts of 8 to 90 wt %, preferably 25 to 75 wt %, most preferably 40 to 70 wt % and
5 C being present in amounts of 0 to 85 wt %, preferably 15 to 65 wt %, most preferably 20 to 50 wt %.

In these blends the fat component B is preferably selected from the group consisting of palm oil; palm oil fractions;
10 cocoa butter equivalents; palm kernel oil; fractions of palm kernel oil; hardened vegetable oils such as hardened palm oil; hardened fractions of palm oil; hardened soybean oil; hardened sunflower oil; hardened rape seed oil; hardened fractions of soybean oil; hardened fractions of
15 rapeseed oil; hardened fractions of sunflower oil; mixtures of one or more of these oils and interesterified mixtures thereof.

Fat component C in general will be a liquid oil and is
20 preferably selected from the group consisting of sunflower oil; olive oil; soybean oil; rape seed oil; palm oil olein; cotton seed oil; olein fractions from vegetable oils and high oleic vegetable oils such as HOSF (=high oleic sunflower oil) or HORP (=high oleic rape seed oil).

25

The blends comprising components A, B and C as disclosed above have excellent properties for application in food products containing a fat phase.

30 The blends can also contain other known micronutrients such as vitamins and minerals. It was however found that it was very beneficial if the blends also contain isoflavones

and/or flavones in amounts corresponding with 0.005 to 5 % of the total amount of ursolic acid and oleanolic acid. In this way compositions are obtained that combine all the health aspects from the ursolic acid and oleanolic acid with the known health aspects from isoflavones respectively flavones (in particular those concerning women's health, in particular for postmenopausal women).

The mixtures and blends according to the invention can be used in food products to provide the health aspects to the food product without giving a negative taste. Therefore part of our invention are also food products with a fat phase comprising at least partly the blend according to the invention and as formulated in claims 3 to 9. The food products can be selected from the group consisting of spreads (fat contents of 10 to 90 wt %); dressings; mayonnaises; cheese; ice creams; ice cream coatings; confectionery coatings; fillings; sauces and culinary product. Very beneficial food products are the food products with a continuous fat phase representing about 10 to 90 wt % of the food product.

According to a last embodiment of our invention our invention also concerns the process for the making of our novel mixtures and blends. Therefore part of our invention also is a process for making a mixture with the composition according to claims 1 and 2 wherein:

- skins of fruit, in particular skins from apples, cranberries, cherries, prunes or pears are extracted with an organic solvent, in particular selected from the group consisting of ketones, esters, alcohols and hydrocarbons

- an extract containing a mixture of ursolic acid and oleanolic acid is isolated
- the organic solvent is removed from the extract and a mixture comprising ursolic acid and oleanolic acid is isolated
- this mixture obtained is dissolved in water or an organic solvent or a mixture thereof, preferably acetone/water in a weight ratio of 50/50 to 95/5 while heating after the mixture is dissolved the solution obtained is cooled to a temperature of maximum 25 °C and crystals formed are separated as the desired product from the rest.

Particulars of our extraction are:

- 15 Prior to the extraction, the fruit skins or herbs are dried to humidity lower than 10 wt%, preferably between 2 and 6 wt%. The skins are then powdered with an electric stirrer. Extraction was performed by mixing the particles with an organic solvent using 1 to 1000 kg of solvent per 1 kg of starting material, preferably 10 to 100 kg of solvent per 1 kg of starting material. The organic solvents are selected from the following groups: alkanes, ketones, esters, aldehydes, hydrocarbons, alcohols. Extraction was performed at temperatures between 0°C and boiling temperature (T_b) of the organic solvent, preferably at temperatures between $T_b - 10$ °C and T_b . The residence time for extraction is not less than 5 minutes, preferably between 15 and 60 minutes. After extraction, the remaining particles were filtered and the solvent was evaporated partly or completely. The remaining extract is washed by dissolving it in organic solvents or water or a mixture of these, at temperatures between 0°C and the boiling temperature of the solvent,

preferably at temperatures above 20 °C. Washing is performed by contacting the extract and washing solvent for at least 15 minutes, preferably between 30 and 60 minutes. The extract is recovered by crystallisation in organic solvents or water or a mixture of these by cooling to temperatures below 25 °C, preferably at 0°C, followed by filtration of the extract.

The blends can be made by a process wherein the mixture according to claim 1 or 2 is mixed in appropriate amounts with a glyceride selected from the glycerides from claims 6 and 7 in amounts appropriate to give the correct amounts according to claim 5, whereupon the total blend is homogenised.

15

EXAMPLES

1. Lab scale extraction

400 g of ground apple skin was mixed with 2 l acetone. The extraction was performed at 58°C for 3 h while stirring. After extraction the solvent was removed and the product dissolved in a mixture of water/acetone (10:90) by heating up to 60°C. Crystallization was performed at room temperature. After 15 min white crystals were formed which were separated by filtration (20 °C). The second crystallization was at 4°C over night. The crystals were separated by filtration and dried under nitrogen. The content of ursolic acid and oleanolic acid was determined with GC by using cholesterol as an internal standard (example 3).

2. Large scale extraction

Extraction was performed in a 180 litre (solvent) crystalliser with a gate-stirrer with hot acetone at 45 - 50°C for about 30 min at ambient pressure. 10 kg
5 ground apple peel/pomace was slurried in 100 l acetone at 35 °C. 10 min was allowed for heating to 45 - 50 °C. After this preheating period, the temperature was kept constant between 45 - 50 °C for about 30 min. After extraction the slurry was led to the A4-filter (pore-size 40 µm) for
10 filtration of the pomace waste. Before feeding the slurry to the filter, the filter was preheated with acetone at 50°C. Filtration was performed by putting up to 1 bar overpressure on the slurry. Filtration of the slurry was performed quickly to prevent crystal formation in the
15 filter cake. The extract of the filtration was sampled to determine the yield and selectivity of the extraction. The evaporation was partly performed in the LUWA evaporator vessel at reduced pressure (50 mbar). Before the product was crystallised from the solvent, the extract product was
20 filtered over a candle filter (10 µm) to remove the last remaining pomace particles. The remaining solvent (20 l) was evaporated in a separate evaporator/boiler. The dried solid extract was washed with demi-water to remove the water-soluble substances. The extract was washed with demi-
25 water for three times, filtrated and dried overnight in an autoclave at 5-10 mbar pressure. The filtrate was washed with 250 ml demiwater, stirring for 0.5 h at 50°C. The mixture was cooled to 0°C for crystallisation of the product. The product was filtrated over a black ribbon
30 paper filter (10 µm) in a Büchner funnel. The sold product was resuspended in 250 ml of hexane and stirred for 0.5 h at 60°C. The mixture was cooled to 0°C for crystallization

of the product. The product was filtrated over a black ribbon paper filter (10 μ m) in a Büchner funnel and dried as described before.

5 3. HPLC analysis

The HPLC analysis to determine the apolar components was performed with the following equipment and conditions:

Apparatus and equipment

10 Analytical balance; Alltech Econoshpere Silica HPLC column (150*4.6 mm, 5 μ m)

Ternair HPLC pump; HPLC autosampler

Alltech 500 ELSD detector (low gain) drift-tube: 75 °C;

Nebulizer: 1.75 L/min

15

HPLC analysis

The gradient that is used for this analysis is described in the appendix. This gradient was run without injecting a sample to flush the column before injecting samples. 5 μ L

20 of the sample solution was injected into the HPLC system.

Gradients used for Silica Straight Phase HPLC

Solvents: A: hexane/toluene (50/50)

B: toluene/ethylacetate/formic acid (500/200/16)

25

Table 1: Determination of diglycerides

time (min)	flow (mL/min)	A	B
initial	0.9	95	5
2	0.9	95	5
3	0.9	95	5
5	0.9	90	10
7	0.9	75	25
8	0.9	50	50
15	0.9	10	90
23	0.9	10	90
31	0.9	95	5
35	0.0	95	5

4. GC analysis

GC analysis to determine the ursolic acid and the low molecular weight components content was performed with the following equipment and conditions:

Injection volume: 0.4 μ L (cold-on-column)

Column: Chrompack SIMDIST (10m*0.53mm, 0.1 μ m filmthickness) or: Quadrex DB-5

10 (10m*0.53mm, 0.1 μ m filmthickness)

Ovenprogram: 1: 120 °C-1 min-20 °C/min, 2: 325 °C-0 min-5.0 °C/min, 3: 350°C-5 min

Detection: FID (360°C)

Carriergas: Hydrogen (column pressure: 15 psi)

15 Internal standard: Cholesterol

5. Taste panel with different extracts

Seven panellists tasted the different extracts.

Objective: Comparison of the taste of the crude extract

20 with the purified extract in a white filling. Reference was the pure white filling.

11

	Intensity score					
	0	1	2	3	4	5
White Filling	7 s					
WF + 2 % AE 1			2 s	1 s	1 s	3 s
WF + 2 % AE 2	1 s	1 s		2 s	3 s	
WF + 2 % PAE 1	4 s	3 s				
WF + 2 % PAE 2	3 s	3 s	1 s			

0 = No taste at all; 1 = A trace of taste; 2 = Faint taste;
 3 = A clear taste; 4 = A strong taste; 5 = A very strong
 taste; WF = White filling; AE = Apple extract; PAE = Pure
 5 apple extract; s = subjects

The majority of the panellists have tasted a clear negative
 taste for the crude extract. The purified extracts however,
 were assessed as tasteless or only had a trace of a taste.

10

CLAIMS

1. Mixture comprising ursolic acid and oleanolic acid in a weight ratio of 1:99 to 99:1, preferably 5:95 to 95:5, most preferably 15:85 to 15:85) wherein the mixture contains less than 20 wt %, preferably less than 10 wt %, most preferably 1 to 6 wt % of the natural apolar and/or low molecular weight components as present in natural extracts for ursolic acid and oleanolic acid.
2. Mixture according to claim 1 wherein the natural apolar and/or low molecular weight components are the components that provide an off taste to the natural extract and belong to the class of hydrocarbons, alcohols, fatty acids, triglycerides, ketones and carbohydrates.
3. A blend of a health component and a glyceride wherein the health component is the mixture according to claims 1 and 2 and which blend contains 1 to 99 wt % of one or more components selected from mono-, di-, and triglycerides as the glyceride.
4. Blend according to claim 3 wherein the glyceride part displays a solid fat content measured by NMR-pulse on a non-stabilised fat at the temperature indicated of :
5 to 90 at 5 oC
2 to 80 at 20 oC and
less than 15, preferably less than 10 at 35 oC
5. Blend according to claim 3 and 4 wherein the blend comprises components A, B and C, wherein:
A = the composition of claims 1 or 2

B = a solid fat with an N20 of more than 20, preferably more than 45, most preferably more than 60 and

C = a fat having at least 40 wt % of fatty acids with 18 C-atoms and having one to three double bonds

A being present in amounts of more than 0.1 wt %, preferably 0.1 to 20 wt %, most preferably 0.2 to 10 wt %

B being present in amounts of 8 to 90 wt %, preferably 25 to 75 wt %, most preferably 40 to 70 wt % and

C being present in amounts of 0 to 85 wt %, preferably 15 to 65 wt %, most preferably 20 to 50 wt %

6. Blend according to claim 5 wherein fat B is selected from the group consisting of palm oil; palm oil fractions; cocoa butter equivalents; palm kernel oil; fractions of palm kernel oil; hardened vegetable oils such as hardened palm oil; hardened fractions of palm oil; hardened soybean oil; hardened sunflower oil; hardened rape seed oil; hardened fractions of soybean oil; hardened fractions of rapeseed oil; hardened fractions of sunflower oil; mixtures of one or more of these oils and intersterified mixtures thereof

7. Blend according to claim 5 wherein fat C is selected from the group consisting of sunflower oil; olive oil; soybean oil; rape seed oil; palm oil olein; cotton seed oil; olein fractions from vegetable oils and high oleic vegetable oils such as HOSF or HORP.

8. Blend according to claim 5, wherein the blend contains a component A that also contains isoflavones and/or

flavones in amounts corresponding with 0.005 to 5 % of the total amount of ursolic acid and oleanolic acid.

9. Blend according to claim 5 wherein component A is a component isolated from fruit skins such as skins from apples, pears, cranberries, cherries or prunes.

10. Food products with a fat phase comprising at least partly the blend according to claims 3 to 9.

11. Food products according to claim 10 wherein the food product is selected from the group consisting of spreads (fat contents of 10 to 90 wt %); dressings; mayonnaises; cheese; ice creams; ice cream coatings; confectionery coatings; fillings; sauces and culinary products.

12. Food products according to claims 10 and 11 wherein the food product comprises 10 to 90 wt % of a continuous fat phase.

13. Process for making a mixture with the composition according to claims 1 and 2 wherein:

- skins of fruit, in particular skins from apples, cranberries, cherries, prunes or pears are extracted with an organic solvent, in particular selected from the group consisting of ketones, esters, alcohols and hydrocarbons.
- an extract containing a mixture of ursolic acid and oleanolic acid is isolated,
- the organic solvent is removed from the extract and a mixture comprising ursolic acid and oleanolic acid is isolated

- this mixture obtained is dissolved in a mixture of water and/or an organic solvent, preferably acetone/water in a weight ratio of 50/50 to 95/5 while heating
- after the mixture is dissolved the solution obtained is cooled to a temperature of maximum 25 °C and crystals formed are separated as the desired product from the rest.

14. Process for making a blend of a glyceride and the mixture according to claims 1 and 2 wherein the mixture according to claim 1 or 2 is mixed in appropriate amounts with a glyceride selected from the glycerides from claims 6 and 7 in amounts appropriate to give the correct amounts according to claim 5, whereupon the total blend is homogenised.

ABSTRACT

Mixtures comprising ursolic acid and oleanolic acid in weight ratios of 1:99 to 99:1 that contain less than 20 wt% of natural apolar components and/or low molecular weight components, normally present in natural extract for ursolic acid and oleanolic acid do not display the negative off taste of the natural extracts anymore.